

# Structure and Enumeration of $K_4$ -minor-free links and link diagrams

Juanjo Rué<sup>1,2</sup>

*Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain*

Dimitrios M. Thilikos<sup>1,2</sup>

*ALGCo project-team, LIRMM, Université de Montpellier, CNRS, Montpellier, France*  
*Department of Mathematics, National and Kapodistrian University of Athens, Athens, Greece*

Vasiliki Velona<sup>1,2</sup>

*Department of Economics, Universitat Pompeu Fabra, Barcelona, Spain*  
*Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain*

---

## Abstract

We study the class  $\mathcal{L}$  of link types that admit a  $K_4$ -minor-free diagram, i.e., they can be projected on the plane so that the resulting graph does not contain any subdivision of  $K_4$ . We prove that  $\mathcal{L}$  is the closure of a subclass of torus links under the operation of connected sum. Using this structural result, we enumerate  $\mathcal{L}$  and subclasses of it, with respect to the minimal number of crossings or edges in a projection of  $L \in \mathcal{L}$ . Further, we enumerate (both exactly and asymptotically) all connected  $K_4$ -minor-free link diagrams, all minimal connected  $K_4$ -minor-free link diagrams, and all  $K_4$ -minor-free diagrams of the unknot.

*Keywords:* series-parallel graphs, links, knots, generating functions, asymptotic enumeration, map enumeration

---

# 1 Introduction

The exhaustive generation of knots and links is a well-established problem in low dimensional geometry (see [7, Ch.5], for instance). However, there are very few enumerative results in the literature and they are relatively recent, (see [10] and [6]). Moreover, there seems to be no known results connecting graph theoretic classes with link classes.

Let us start with some formal definitions. A *knot*  $K$  is a smooth embedding of the 1-dimensional sphere  $\mathbb{S}^1$  in  $\mathbb{R}^3$ . A *link* is a finite disjoint union of knots  $L = K_1 \cup \dots \cup K_n$ . Two links  $L_1$  and  $L_2$  are equivalent if there is a continuous and injective function  $h : \mathbb{R}^3 \times [0, 1] \rightarrow \mathbb{R}^3$ , such that  $h(L_1, 0) = L_1$  and  $h(L_1, 1) = L_2$ . A link equivalent to a set of non intertwined circles is called a *trivial link*. If it is a knot, we also call it the *unknot*.

Consider a link  $L$  and a sphere  $\mathbb{S}^2$  embedded in such a way that it meets the link transversely in exactly two points  $P_1$  and  $P_2$ . Then we can discern two different links  $L_1, L_2$ , after connecting  $P_1$  and  $P_2$ . The first corresponds to the part of  $L$  in the interior of the sphere and the second to the part in the exterior. We then call  $L$  a *connected sum* with factors  $L_1, L_2$ , denoted  $L_1 \# L_2$ . A link that does not have non-trivial factors is called *prime*, otherwise *composite*. A link is *split* if there is a sphere embedded in the link complement that separates the link. Each of the components is called a *disjoint component* of the link and, conversely, the link is their *disjoint sum*.

Let  $L \in \mathbb{R}^3$  be a link and let  $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  be a projection map. If for all  $x \in L$ ,  $|\pi^{-1}(x)| = 2$  or  $1$ , and all the double points are finite and transverse, then the projection is said to be *regular*. For all knots defined here, there exists a regular projection [2, Ch. 3]. This allows to work with *link-diagrams*, i.e. a triple  $(V, E, \sigma)$ , where  $(V, E)$  is a 4-regular plane graph and  $\sigma : V(G) \rightarrow \binom{E(G)}{2}$ , where, for every  $v \in V(G)$ ,  $\sigma(v)$  is a set of two opposite edges of the embedding of  $G$  and encodes which pair is overcrossing. Notice that a link-diagram may also have vertexless edges. We call these edges *trivial*. The *crossing number* of a link  $L$  is the minimum number of crossings that can have a diagram of it. Such a diagram is called *minimal*. We also say that a diagram  $(V, E, \sigma)$  is  $K_4$ -minor-free if the graph  $(V, E)$  does not contain

---

<sup>1</sup>The first and the third author have partially been supported by the MINECO projects MTM2014-54745-P, MTM2017-82166-P and the María de Maetzu research grant MDM-2014-0445 (BGSMath). The second author has been supported by the project “ESIGMA” (ANR-17-CE40-0028). The third author has been supported under an FPI grant from the MINECO research project MTM2015-67304-PI.

<sup>2</sup>Emails: [juan.jose.rue@upc.edu](mailto:juan.jose.rue@upc.edu), [sedthilk@thilikos.info](mailto:sedthilk@thilikos.info), [vasiliki.velona@upf.edu](mailto:vasiliki.velona@upf.edu)

any subgraph that is a subdivision of a  $K_4$ .

A *torus link* is a link that can be embedded on the torus. They are denoted by  $T(p, q)$ ,  $p, q \in \mathbb{Z}$ , where  $p$  and  $q$  are the number of times that the link crosses the meridian and the longitude cycle, respectively. Let  $\mathcal{T}_2$  be the closure of torus links of type  $T(2, q)$ ,  $q \in \mathbb{Z} - \{0\}$ , under the connected sum operation. Let  $\mathcal{L}$  be the class of links that have a  $K_4$ -minor-free link-diagram and  $\mathbf{mcl}(\mathcal{T}_2)$  be the closure of  $\mathcal{T}_2$  under disjoint sum.

## 2 Structure of $K_4$ -links and enumeration

Our first result gives the type of links that admit a  $K_4$ -minor-free diagram.

**Theorem 2.1** *The links that admit a  $K_4$ -minor-free diagram are exactly the ones in  $\mathbf{mcl}(\mathcal{T}_2)$ , i.e.  $\mathcal{L} = \mathbf{mcl}(\mathcal{T}_2)$ .*

Based on the above, we derive a precise description of links that admit a  $K_4$ -minor-free diagram. Let  $\bar{\mathcal{L}}$  be the class of non-split links in  $\mathcal{L}$  and  $\hat{\mathcal{L}}$  the class of links in  $\mathcal{L}$  with no trivial disjoint components, with size defined as their crossing number. We will obtain the asymptotic growth of  $\mathcal{L}$  with respect to the number of edges in a minimal diagram (not the crossing number, so as to account also for trivial disjoint components). Let  $L(z)$ ,  $\bar{L}(z)$ , and  $\hat{L}(z)$  be the respective generating functions. Finally, let  $\mathcal{T}$  be the class of non plane, unrooted trees, where the vertices are labelled with multisets of the set  $\{1\} \cup \{\pm 3, \pm 5, \dots\}$  and the edges are labelled with an element of the set  $\{2, \pm 4, \pm 6, \dots\}$ . These labels will encode crossing numbers, hence a label  $i$  corresponds to size  $|i|$ . The size of  $T \in \mathcal{T}$  is the sum of its labels. See Figure 1 for an example of such a tree.




Figure 1. An object of  $\mathcal{T}$ .

**Proposition 2.2**  $\bar{\mathcal{L}} \cong \mathcal{T}$  and  $\mathbf{MSet}_{\geq 1}(\mathcal{T} \setminus T_1) \cong \hat{\mathcal{L}}$ , where  $\mathbf{MSet}$  is the multiset operator with at least one element and  $T_1 \in \mathcal{T}$  is the tree with one vertex and label 1.

Using Proposition 2.2 and the framework of the *Symbolic Method* (see [4]), we derive combinatorial expressions for  $\mathcal{L}$  that translate to generating functions. Let  $\bar{\mathcal{K}}$  and  $\mathcal{K}$  be the classes of prime knots and composite knots plus the trivial knot in  $\mathcal{L}$ , respectively. For prime knots in  $\mathcal{L}$ , i.e. prime torus knots  $T(2, 2i + 1)$ ,  $i \in \mathbb{Z} \setminus \{0, -1\}$ , it holds that  $\bar{K}(z) = 2 \sum_{i \geq 1} z^{2i+1} = \frac{2z^3}{1-z^2}$ , counting according to their crossing number. For  $\mathcal{K}$ , it is enough to consider non-empty multisets of prime torus knots, therefore  $\mathcal{K} = \text{MSet}_{\geq 1}(\bar{\mathcal{K}})$ , which translates to the expression

$$K(z) = \exp \left( \sum_{k \geq 1} \frac{1}{k} \bar{K}(z^k) \right) - 1 = \exp \left( \sum_{k \geq 1} \frac{1}{k} \frac{2z^{3k}}{1-z^{2k}} \right) - 1.$$

Let  $\mathcal{G}$  be the class of unrooted and unlabelled non-plane trees. Notice that one cannot replace immediately  $z$  for  $\mathcal{K}$  in the ogf  $G(z)$ , since the vertices are not distinguishable. Hence, to continue, one needs to use cycle indices. Let  $\mathcal{G}^\bullet$  be the class of rooted and unlabelled non-plane trees. For the cycle index of  $\mathcal{G}^\bullet$ , it is known (see [1]) that

$$\mathcal{Z}_{\mathcal{G}^\bullet}(s_1, s_2, \dots) = s_1 \exp \left( \sum_{k \geq 1} \frac{1}{k} \mathcal{Z}_{\mathcal{G}^\bullet}(s_k, s_{2k}, \dots) \right).$$

Let  $\mathcal{E}$  be the combinatorial class of the edge labels of  $\mathcal{T}$ , hence  $E(z) = z^2 + \frac{2z^4}{1-z^2}$ , and let  $f(z) = E(z)K(z)$ . We can now obtain the ordinary generating function of  $\mathcal{F} = \mathcal{T}^\bullet \circ (\mathcal{E} \times \mathcal{L}_c)$ . By Polya's Theorem, the latter satisfies the equation  $F(z) = \mathcal{Z}_{\mathcal{G}^\bullet}(f(z), f(z^2), \dots)$ . Then, it holds that

$$\bar{L}(z) = \frac{F(z)}{E(z)} + \frac{E(z)}{2} \left( -\frac{F(z)^2}{E(z)^2} + \frac{F(z^2)}{E(z^2)} \right),$$

by the Dissymmetry Theorem (see [1]). The first terms of  $\bar{L}(z)$  are as follows:

$$\bar{L} = 1 + z^2 + 2z^3 + 3z^4 + 4z^5 + 9z^6 + 12z^7 + 26z^8 + 40z^9 + \dots$$

We can then get asymptotic estimates by means of complex analytic tools:

**Theorem 2.3**  $\bar{\mathcal{L}}$  has asymptotic growth of the form:

$$[z^n] \bar{L}(z) \sim \frac{cn^{-3/2}}{\Gamma(-1/2)} \rho^{-n}, \quad \rho \approx 0.44074, \quad c \approx 1.45557.$$

Additionally,  $\hat{\mathcal{L}}$  has the same type of asymptotic growth with the same  $\rho$  and  $c \approx 3.61691$ .

The generating function  $L(z)$  is equal to  $\hat{L}(z^2) \frac{1}{1-z}$ , since a link diagram of  $n$  vertices has  $2n$  non-trivial edges and a number of trivial edges. We thus obtain the following corollary.

**Corollary 2.4**  $\mathcal{L}$  has asymptotic growth of the form:

$$[z^n]L(z) \sim \frac{cn^{-3/2}}{\Gamma(-1/2)}\rho^{-n},$$

where  $\rho \approx 0.44074$  and  $c \approx 8.97779$  or  $c \approx 3.95687$ , depending on whether  $n$  is even or odd, respectively.

Finally, we also get asymptotic estimates for the coefficients of  $K(z)$ . The following result is a consequence of Meinardus' Theorem (see [4, VIII.23]).

**Theorem 2.5**  $\mathcal{K}$  has asymptotic growth of the form:

$$[z^n]K(z) \sim cn^{-7/4} \exp(c'n^{1/2}),$$

where  $c = \frac{e^{2 \ln(2)\zeta(0)}(\Gamma(2)\zeta(2))^{5/4}}{2\sqrt{\pi}} \approx 0.26275$ ,  $c' = 2\sqrt{\Gamma(2)\zeta(2)} \approx 2.56509$ , and  $\zeta(z)$  is the Riemann zeta function.

### 3 Enumeration of families of link diagrams

Enumerating connected  $K_4$ -free link-diagrams is equivalent to enumerating 4-regular unrooted planar maps which are  $K_4$ -minor-free, that we call  $\bar{\mathcal{M}}_1$ . We first give a construction for the rooted ones,  $\mathcal{M}_1$ , with respect to edges, adapting the construction for rooted 4-regular maps in [8]. With this we obtain a functional system of equations that can be analyzed by using the analytic machinery developed by Drmota in [3]. Finally, by adapting an argument by Richmond and Wormald in [9], we are able to show that we can unroot the maps under study, showing that the maps in our class with non-trivial automorphisms are exponentially few. This study gives the following result:

**Theorem 3.1** The class of connected  $K_4$ -free link diagrams  $\bar{\mathcal{M}}_1$  satisfies that

$$[z^n]\bar{\mathcal{M}}_1(z) \sim \frac{1}{2n} \frac{cn^{-3/2}}{\Gamma(-1/2)}\rho^{-n}2^n, \quad \rho \approx 0.31184, \quad c \approx 1.52265$$

Refining the combinatoric analysis in the previous case (with more auxiliary classes) and using again an adaptation of the argument of Richmond and Wormald in [9], we are able to asymptotically enumerate also the class of minimal link diagrams  $\bar{\mathcal{M}}_2$ , as well as the class of  $K_4$ -free link diagrams of the unknot,  $\bar{\mathcal{M}}_3$ :

**Theorem 3.2** The class of connected  $K_4$ -free minimal link diagrams  $\bar{\mathcal{M}}_2$ , and

the class of  $K_4$ -free link diagrams of the unknot,  $\bar{M}_3$  satisfy:

$$[z^n]\bar{M}_2(z) \sim \frac{1}{2n} \frac{c_2 n^{-3/2}}{\Gamma(-1/2)} \rho_2^{-n}, \quad \rho_2 \approx 0.41456, \quad c_2 \approx 0.81415,$$

$$[z^n]\bar{M}_3(z) \sim \frac{1}{2n} \frac{c_3 n^{-3/2}}{\Gamma(-1/2)} \rho_3^{-n}, \quad \rho_3 \approx 0.23188, \quad c_3 \approx 2.19020.$$

**Acknowledgement.** All authors like to thank Lefteris Kirousis that contributed to the inception of this research.

## References

- [1] F. Bergeron, G. Labelle, and P. Leroux. *Combinatorial species and tree-like structures*, volume 67. Cambridge University Press, 1998.
- [2] P. R. Cromwell. *Knots and links*. Cambridge University Press, 2004.
- [3] M. Drmota. *Systems of functional equations. Random Structures and Algorithms*, 10, 103-124, 1997.
- [4] P. Flajolet and R. Sedgewick. *Analytic combinatorics*. Cambridge University press, 2009.
- [5] A. Kawauchi. *A survey of knot theory*. Birkhäuser, 2012.
- [6] S. Kunz-Jacques and G. Schaeffer. The asymptotic number of prime alternating links. In *FPSAC'2001*, pages 10–p. Arizona State University, 2001.
- [7] W. Menasco and M. Thistlethwaite. *Handbook of knot theory*. Elsevier, 2005.
- [8] M. Noy, C. Requilé, and J. Rué. Enumeration of labeled 4-regular planar graphs. *Submitted*, available on-line at [arXiv: 1709.04678](https://arxiv.org/abs/1709.04678).
- [9] L. B. Richmond and N. C. Wormald. Almost all maps are asymmetric. *J. of Combinatorial Theory, Series B*, 63(1):1–7, 1995.
- [10] C. Sundberg and M. Thistlethwaite. The rate of growth of the number of prime alternating links and tangles. *Pacific J. of mathematics*, 182(2):329–358, 1998.